

ISSN : 2309-3234

TMSS MEDICAL COLLEGE JOURNAL

Volume 05, No. 02, July 2015



An Official Publication of TMSS Medical College, Bogra



TMSS Medical College Journal

July 2015

An Official Publication
of
TMSS Medical College, Bogra

TMSS Medical College Journal

Vol. 05, No. 02, July 2015

An Official Publication of TMSS Medical College, Bogra, Bangladesh

Editorial Board

Chairperson : Prof. Dr. Most. Rebeca Khatoun
Editor-In-Chief : Prof. Dr. Md. Abdus Shukur
Executive Editor : Prof. Dr. Md. Azfarul Habib
Publication Editor : Dr. Md. Abdul Mazid
Finance Editor : Dr. Md. Matiur Rahman

Members

Prof. Dr. M. A. Gafur
Prof. Dr. Md. Aynul Huq
Prof. Dr. Md. Shahjahan Ali Sarkar
Prof. Dr. Nur-A-Farhana Islam
Dr. Md. Azizar Rahman
Dr. Md. Sheik Abdul Muktaf
Dr. Md. Kamrul Islam

Review Committee

Prof. Dr. Raghib Ahsan
Prof. Dr. Md. Anup Rahman Chowdhury
Prof. Dr. Md. Abul Hossain Khan

Advisory Board

Prof. Dr. Hosne-Ara Begum
Prof. Dr. A.K.M Ahsan Habib
Prof. Dr. Syed Mainul Hasan
Prof. Dr. S.A.M. Israrul Bari
Prof. Dr. Md. Rafiqul Islam
Prof. Dr. Latifur Rahman
Prof. Dr. Abdul Gaffar
Prof. Dr. Hafiza Arzuman
Dr. A.K.M. Ahsan Habib
Dr. Md. Mostafa Alam

Published by

Dr. Md. Abdul Mazid
on behalf of the TMSS Medical College

Graphic Designer

Md. Sirajul Islam
TMSS, Bogra

Printed by

TMSS Printing Press
Bogra, Bangladesh

The TMSS Medical College Journal is a peer reviewed Journal. It is published two times a year, (January and July). It accepts original articles, review articles and case reports. Complimentary copies of the journal are sent to libraries of all medical and other relevant academic institutions in the country and selected institutions abroad.

While every effort is always made by the Editorial Board and the members of the Journal Committee to avoid inaccurate or misleading information appearing in the TMSS Medical College Journal, information within the individual article are the responsibility of its author(s). The TMSS Medical College Journal, its Editorial Board and Journal Committee accept no liability whatsoever for the consequences of any such inaccurate and misleading information, opinion or statement.

Address of Correspondence

Executive Editor, TMSS Medical College Journal
TMSS Medical College, Rangpur Road
Thengamara, Bogra, Bangladesh.
Phone: 051-61830, Fax : 051-61830
e-mail: editortmcjournal@gmail.com
Web: www.tmssmedicalcollege.com

Association of Serum Calcium with Acute Myocardial Infarction

Chowdhury SN^{1*}, Sultana N², Chowdhury AH³, Mazumder P⁴, Rahman T⁵, Sharmin N⁶

1. Dr. Sanjeela Nahreen Chowdhury, Assistant professor, Department of Biochemistry, Universal Medical College, Dhaka.
2. Dr. Nasima Sultana, Professor & Head, Department of Biochemistry, Dhaka Medical College, Dhaka.
3. Dr. Amirul Hossain Chowdhury, Head, Department of Forensic Medicine, Shaheed M. Monsur Ali Medical College, Sirajgonj.
4. Dr. Pinke Mazumder, Assistant professor, Department of Biochemistry, Army Medical college, Bogra.
5. Dr. Taniza Rahman, Assistant professor, Department of Biochemistry, Ashiyan Medical college, Dhaka
6. Dr. Nazia Sharmin, Assistant professor, Department of Biochemistry, Monno Medical college, Manikganj.

* Corresponding Author

Abstract

Background: Acute myocardial infarction (AMI) is a leading cause of morbidity and mortality worldwide. The incidence of AMI is dependent on certain traditional predisposing risk factors. In addition with the traditional risk factors, raised serum calcium level is also being considered as an associated factor for AMI. **Objective:** The aim of the study is to evaluate the association of serum calcium level with acute myocardial infarction in a tertiary level hospital in Bangladesh. **Materials and Methods:** A case control study was conducted in the Department of Biochemistry, Dhaka Medical College, Dhaka from January 2013 to December 2013. In this study, 50 diagnosed cases of AMI from DMCH and 50 age and sex matched apparently healthy subjects were selected from DMCH purposively according to the selection criteria. Blood pressure, height and weight were measured and BMI was calculated. Biochemical parameters-serum calcium and fasting blood glucose in two groups were estimated in mmol/L respectively. Then serum calcium was compared between two groups to observe the association with AMI. Results were analyzed statistically in SPSS version 17.0. Unpaired student's t-test and Spearman correlation analysis were done. All the results were expressed as mean $\pm$ SD and p value <0.05 was accepted as level of significance. **Results:** Present study showed that serum calcium level was significantly higher in cases when compared with that of controls ($p=0.001$) which was 2.61 ± 0.15 mmol/L and 2.13 ± 0.15 mmol/L in cases and controls respectively. Serum calcium also showed a significant positive correlation with AMI ($\rho=0.858$, $p=0.001$). **Conclusion:** Findings of study concludes that increased serum calcium is associated with AMI in a tertiary level hospital in Bangladesh.

Key words: Acute myocardial infarction, Serum calcium, Hypercalcemia.

Introduction

The calcium ion is an essential regulator in many homeostatic systems, including vascular tone, hormone secretion, and intermediary metabolism (Campbell et al., 1983)¹. Calcium is also essential for nerves, muscles, heart and other body system to work properly.

Calcium ion not only plays a role in regulating variety of physiological events, it is also involved in developing pathological condition mainly in heart. Increased level of calcium causes coronary artery and peripheral arteriolar constriction by binding with the heart and smooth muscle through calcium receptor and thus increases cardiac contractility and causes diminished oxygen supply

to myocardium. This eventually increases the risk of cardiovascular disease. According to Nadia (2011)², increased level of calcium causing abnormality of heart leading to excessively forceful or tight contraction causes increased and irregular heartbeat (tachycardia and arrhythmia).

A calcium overload also causes increased blood pressure, coronary artery calcification^{3, 4, 5} and progression of atherosclerosis that involves lipid as well as collagen, elastin and calcium accumulation in coronary vasculature. Myocyte infiltration, endothelial injury, smooth muscle proliferation and migration are also involved in coronary atherosclerosis. Several of these processes are also mediated by calcium ion⁶.

Moreover, calcium is involved in other events such as coronary spasm, thrombus formation and disruption of atherosclerotic plaque, occlusion of coronary artery following rupture of vulnerable atherosclerotic plaque which can cause myocardial infarction and sudden death⁷. Various studies in different parts of the world suggest that serum calcium is strongly associated with myocardial infarction^{8, 9}. In a study Lind et al. (1997)⁹, serum calcium has been found to be an independent, prospective risk factor for myocardial infarction in Swedish middle-aged men. In another study by Palmer et al. (1987)¹⁰ suggested that not only hypercalcemia is associated with AMI, but also slightly elevated serum calcium levels within the normal range are of importance.

The incidence of myocardial infarction is also increasing in Bangladesh. Rates of cardiovascular disease have risen greatly in low-income and middle-income countries^{11, 12, 13} with about 80% of the burden now occurring in these countries. Effective prevention needs a global strategy based on knowledge of the importance of risk factors for cardiovascular disease in different geographic regions and among various ethnic groups. Risk factors for coronary heart disease vary between populations, e.g., lipids are not associated with this disorder in South Asians¹³ and increases in blood pressure might be more important in Chinese people.¹⁴ Known risk factors (generally smoking, hypertension, raised lipids, and diabetes) are being identified and treated. But the incidence rate is still high because these risk factors have been claimed to account for only about half the risk of a myocardial infarction¹³. So, there is a pressing need to identify the associated risk factors of AMI to reduce the incidence and mortality rate.

The relationship between serum calcium and AMI is becoming clear day by day from different studies done around the world. But very little

study is reported so far in Bangladesh. Thus, the present study was designed to observe the association of serum calcium concentration with acute myocardial infarction in Bangladeshi population.

Materials and Methods

A case control study was conducted in the Department of Biochemistry, Dhaka Medical College, Dhaka from January 2013 to December 2013. In this study 50 diagnosed cases of AMI and 50 age and sex matched apparently healthy subjects were selected purposively according to selection criteria. The cases were selected from DMCH. The diagnosis of AMI was based on the electrocardiogram, ischemic cardiac pain lasting at least 30 min, and change in Troponin I. Variables such as smoking, history of hypertension, diabetes mellitus, hypercholesterolemia, height, body weight, current medication prior to the AMI during admission were recorded. Blood pressure, height and weight were measured and BMI was calculated. Biochemical parameters- fasting serum glucose and serum calcium in two groups were estimated in mmol/L respectively. BMI, blood pressure, serum calcium and fasting serum glucose were compared between cases and controls. Results were analyzed statistically in SPSS version 17.0. Unpaired student's t-test and Spearman correlation were done. All the results were expressed as mean $\pm$ SD and p value <0.05 was accepted as level of significance.

Results

Demographic characteristics of the patients are presented in table I. All general characteristics are significantly different between cases and controls except for ages and blood glucose. Mean ($\pm$ SD) of age (in years) of the cases and controls were 53.30 ± 6.74 and 51.86 ± 7.30 respectively. There was no statistically significant difference of mean age between two group ($p=0.308$).

Table I: Demographic characteristics of the subjects

Variables	Case n=50 Mean±SD	Control n=50 Mean±SD	p value
Age (years)	53.30 ± 6.74	51.86 ± 7.30	0.308
Male sex n (%)	31 (64%)	31 (64%)	0.001
Female sex n (%)	19(38%)	19(38%)	
SBP (mm of Hg)	137.80±14.92	124.70±11.35	
DBP (mm of Hg)	88.70±10.24	79.60±7.61	
BMI(Kg/m ²)	27.34±3.37	25.30±3.42	0.001

Unpaired students t-test was done to measure the level of significance; Significance = (p<0.05).

Table I also showed that mean of SBP was significantly higher in cases when compared with that of controls (p =0.001) which were 137.80±14.92 mm of Hg and 124.70±11.35 mm of Hg respectively. Mean (±SD) of DBP was also significantly higher in cases than that of controls (p=0.001).

Table II: Biochemical parameters of study subjects

Variable	Case (n=50) Mean±SD	Control (n=50) Mean±SD	P value
Serum calcium (mmol/ L)	2.61 ± 0.16	2.13 ± 0.15	0.001
Fasting blood sugar (mmol/L)	5.3±0.821	5.2±0.709	0.297

Unpaired student's t-test was done to measure the level of significance. Significant= (P<0.05).

Table II shows the biochemical parameters of the study subjects. Mean (±SD) of serum calcium were 2.61±0.16 mmol/L and 2.13±0.15 mmol/L in cases and controls respectively and it was significantly higher in cases (p=0.001). Fasting blood glucose of the cases and controls were 5.3±.821 and 5.2±.709 respectively. There was no statistically significant difference of mean blood glucose between two group (p=0.297).

Table III: Correlation between s. calcium and AMI (yes/no).

Study subjects	rho value	p value
AMI (Y/N)	0.858	0.001

rho= Spearman's correlation coefficient; Significant= (P<0.05).

The correlation of serum calcium and AMI was done by Spearman correlation test. The result shows significant positive correlation (rho=.858, p=.001) between serum calcium and AMI (Table III).

Discussion

In the present study, Serum glucose was estimated in cases and controls to exclude diabetes. All the subjects in the study were non diabetic with the mean (±SD) fasting serum glucose 5.3±8.21 mmol/L and 5.2±.709 mmol/L in cases and controls respectively. Serum Troponin I was estimated and ECG monitoring was done to diagnose the AMI.

The present study revealed that the mean (±SD) serum calcium concentration was 2.61±.16 mmol/L and 2.13 ± 0.15 mmol/L in AMI cases and healthy controls respectively. In the current study we found that, serum calcium level was significantly higher (p=.001) in AMI cases than that of controls. Similarly, previous study by Lind et al. (1997)⁹ found that subjects with a history of MI had significantly higher serum calcium than those without infarction. Our finding of a positive association between elevated serum calcium level and AMI is also consistent with other studies^{8, 15}. Herrmann et al. (1986)¹⁶ demonstrated that increased serum calcium predisposes calcific deposition in the valve cusps and coronary arteries, which could cause significant aortic valve stenosis and accelerate coronary atherosclerosis, this can mismatch the myocardial oxygen supply and demand.

The mechanisms underlying the associations between circulating calcium, cardiovascular risk factors and cardiovascular disease appear to be multiple and complex. The calcium-sensing receptors are expressed in the vascular smooth muscle and endothelial cells and mediate some of the effect of circulating calcium on vascular tone¹⁷. Thus, serum calcium may be involved in regulating blood pressure by controlling vascular smooth muscle cell contractility and modulating peripheral vascular resistance^{18, 19}.

The present study showed mean ($\pm$ SD) of SBP and DBP in cases were 137.8 ± 14.92 mm of Hg and 88.70 ± 10.24 mm of Hg and in controls 124.7 ± 11.35 mm of Hg and 79.60 ± 7.61 mm of Hg respectively. This findings supports the study in a Swedish population where it has shown that the mean value of SBP and DBP were 138 ± 21 mm of Hg and 87 ± 13 mm of Hg in cases and 132 ± 17 mm of Hg and 83 ± 10 mm of Hg in controls respectively⁹. Current study showed that the difference was statistically significant between two groups ($p=0.001$) in respect of SBP and DBP. In contrast to our findings, the study by Buckley (1987)²⁰ in 325 males and the study by Andersen (1984)²¹ in 70 men and women from Denmark have reported hypertension is associated with lower serum calcium. These inconsistencies might have occurred due to different selection criteria used for study subjects and population, and different methodology.

In this study, mean ($\pm$ SD) of BMI was 27.34 ± 3.37 Kg/m² in AMI cases which was significantly higher than that of controls ($p=.001$). In this study subjects with AMI has higher BMI and higher serum calcium than that of controls, similar to other studies^{9, 22}

In our study, Spearman correlation coefficient shows a significant positive correlation between serum calcium and AMI subjects ($\rho=0.858$, $p=.001$) suggesting increased serum calcium level

significantly increases the chances of AMI. Similarly, Cohort study by John et al. (2013)²³ suggested that plasma calcium is a predictor of CVD and a predictor of MI^{8, 9} as well as a predictor of cardiovascular mortality^{24, 25}. However, in Framingham study and Atherosclerosis risk in Communities (ARIC) study calcium was not found to be a predictor of cardiovascular disease in age- and sex-adjusted or multivariable-adjusted models^{26, 27}. Another study done by Jin Y et al. (2013)²⁸ demonstrated that serum calcium levels are not associated with IHD. The conflicting results from these studies may reflect demographic differences between participants or differences in analysis. Findings of our study suggested that, increased serum calcium is positively associated with AMI in both sexes.

Conclusion

This study permits to conclude that increase serum calcium is associated with AMI. However, the combined evaluation of increased serum calcium and other risk factors might help to assess the risk of future occurrence of acute myocardial infarction as well as reduce the incidence of AMI.

References

1. Campbell, A. K. 'Intracellular Calcium, its universal role as regulator', Wiley, USA 1983.
2. Nadia, H. 'Calcium and cardiac function', Journal of Biomedical science 2011; 20: pp. 214.
3. Rubin, M.R., Rundek, T., Mc Mohan, D.J., Lee, H.S. and Silverbergh, S.J. 'Carotid artery plaque thickness is associated with increased serum calcium levels: the Northern Manhattan study'. Atherosclerosis 2007; 194(2): pp.426-32.

4. Bolland, M.J., Avenell, A., Baron, J.A., Grey, A., MacLennan, G.S., Gamble G.D. and Reid I.R. 'Effect of calcium supplements on risk of myocardial infarction and cardiovascular events: meta-analysis', *BMJ* 2010; 341(3691): pp. 341.
5. Mark, B., Andrew, G. and Ian, R. 'Calcium and cardiovascular risks', *Australian prescriber* 2013; 36(5): 148-9.
6. Nayler, W.G. 'Calcium, calcium antagonism, atherosclerosis and ischemia', *Journal of cardiovascular Pharmacology* 1992; 19 (2): S17-S21.
7. Schmitz, G., Hankowitz, J., and Kovacs, E.M. 'Cellular processes in atherogenesis: Possible targets of Calcium channel blockers', *Atherosclerosis* 1991; 88(2-3):109-132
8. Jorde R., Johan, S., Patrick, F. and Kaare, H.B. 'Serum calcium and cardiovascular risk factors and diseases; The Tromso study', *Hypertension* 1999; 34: pp. 484-490.
9. Lind, L., Skarfors, E., Berglund, L., Lithell, H. and Ljunghall, S. 'Serum calcium: a new, independent, prospective risk factor for myocardial infarction in middle aged men followed for 18 years', *J clin epidemiol*, 1997; 50(8): pp. 967-973.
10. Palmer, M., Adami, H.O., Bergstrom, R., Jakobsson, S., Akerstrom, G. and Ljunghall, S. 'Survival and renal function in persons with untreated hypercalcaemia: A population-based cohort study with 14 years of follow-up', *The Lancet* 1987; 1(8524): pp. 59-62.
11. Yusuf, S., Reddy, S., Ôunpuu, S., and Anand, S. 'Global burden of cardiovascular diseases, part I: general considerations, the epidemiologic transition, risk factors, and impact of urbanization', *Circulation* 2001; 104: 2746-2753
12. Pais, P., Pogue, J., Gerstein, H. 'Risk factors for acute myocardial infarction in Indians: a casecontrol study', *Lancet* 1996; 348: 358-363.
13. Yusuf, S., Reddy, S., Ôunpuu, S., and Anand, S. 'Global burden of cardiovascular diseases, part II: variations in cardiovascular disease by specific ethnic groups and geographic regions and prevention strategies', *Circulation* 2001; 104: 2855-2864
14. Canto, J.G. and Iskandrian, A.E. 'Major risk factors for cardiovascular disease: debunking the "only 50%" myth', *JAMA* 2003; 290: 947-949
15. Sabanayagam, C., and Shankar, A. 'Serum calcium levels and hypertension among US adults', *J Clin Hypertens* 2011; 13(10): pp. 716-721.
16. Hermann, G., Hehrmann, R., Scholz, H.C., Atkinson, P.L. and Muhlen, V.A. 'Parathyroid hormone in coronary artery disease—results of a prospective study', *J Endocrinol Inves* 1986; 9(4): pp. 265-271.
17. Ward, B.K., Magno, A.L., Walsh, J.P. and Ratajczak, T. 'The role of the calcium-sensing receptor r in human disease', *Clin Biochem* 2012; 45(12): pp. 943-953.
18. Zemel, M.B. 'Regulation of adiposity and obesity risk by dietary calcium: mechanisms and implications', *J Am Coll Nutr* 2002; 21(2): pp. 146S-151S
19. Zemel, M.B. 'Nutritional and endocrine modulation of intracellular calcium: implications in obesity, insulin resistance and hypertension' *Mol Cell Biochem* 1998; 188(1-2): pp. 129-136.
20. Buckley, B.M., Smith, S.C., Beevers, M., Beevers, D.G., McKiernan, M.J. 'Lack of evidence of low ionized calcium levels in

- systemic hypertension' *Am J Cardiol* 1987; 59(8), pp. 878–880.
21. Andersen, F. N., Hedegaard, L., and Thode, J. 'Sex-dependent relation between ionized calcium in serum and blood pressure' *Clin Chem* 1984; 30(1): pp. 116–118.
 22. Farhangi, M.A., Ostadrahimi, A., Mahboob, S. 'Serum calcium, magnesium, phosphorus and lipid profile in healthy Iranian premenopausal women' *Biochemia Medica* 2011; 21(3): 312-20.
 23. John, P.W., Mark, L.D. and Mathew, W.K. 'Plasma calcium as a predictor of cardiovascular disease in a community based cohort' *Clin Endocrinol*, 2013; 78(6): pp. 852-857.
 24. Leifsson, B.G. and Ahrén, B. 'Serum calcium and survival in a large health screening program', *J Clin Endocrinol Metab*, 1996; 81(6): pp. 2149-2153.
 25. Larsson, T.E., Olauson, H., Hagstrom, E., Ingelsson, E. and Arnio, J. 'Conjoint effects of serum calcium and phosphate on risk of total, cardiovascular, and non-cardiovascular mortality in the community', *Arterioscler Thromb Vasc Biol*,
 26. Dhingra, R., Sullivan, L.M., Fox, C.S., Wang, T. J., D'Agostino, R. B. and Gaziano, M. et al. 'Relations of serum phosphorus and calcium levels to the incidence of cardiovascular disease in the community', *Arch Intern Med* 2007; 167: pp. 879–885.
 27. Foley, R.N., Collins, A.J., Ishani, A. and Kalra, P.A. 'Calcium-phosphate levels and cardiovascular disease in community-dwelling adults: the Atherosclerosis Risk in Communities (ARIC) Study', *Am Heart J* 2008; 156 (3): pp. 556–563.
 28. Jin, Y., He, L., Chen, Y., Ren, X., Tang, H. et al. 'Serum calcium levels are not associated with coronary heart disease', *Vasc Health Risk Manag* 2013; 9: 517-20.